

PARENT & ATHLETE CONCUSSION INFORMATION SHEET



WHAT IS A CONCUSSION?

A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by a bump, blow, or jolt to the head or body that causes the head and brain to move quickly back and forth. Even a "ding," "getting your bell rung," or what seems to be a mild bump or blow to the head can be serious.

WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury.

If an athlete reports one or more symptoms of concussion after a bump, blow, or jolt to the head or body, s/he should be kept out of play the day of the injury. The athlete should only return to play with permission from a health care professional experienced in evaluating for concussion.

DID YOU KNOW?

- Most concussions occur without loss of consciousness.
- Athletes who have, at any point in their lives, had a concussion have an increased risk for another concussion.
- Young children and teens are more likely to get a concussion and take longer to recover than adults.

SYMPTOMS REPORTED BY ATHLETE:

- Headache or "pressure" in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not "feeling right" or is "feeling down"

SIGNS OBSERVED BY COACHING STAFF:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes
- Can't recall events prior to hit or fall
- Can't recall events after hit or fall

▶ **"IT'S BETTER TO MISS ONE GAME THAN THE WHOLE SEASON"**

CONCUSSION DANGER SIGNS

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. An athlete should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that gets worse
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless, or agitated
- Has unusual behavior
- Loses consciousness (even a brief loss of consciousness should be taken seriously)

WHAT SHOULD YOU DO IF YOU THINK YOUR ATHLETE HAS A CONCUSSION?

1. If you suspect that an athlete has a concussion, remove the athlete from play and seek medical attention. Do not try to judge the severity of the injury yourself. Keep the athlete out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.
2. Rest is key to helping an athlete recover from a concussion. Exercising or activities that involve a lot of concentration, such as studying, working on the computer, and playing video games, may cause concussion symptoms to reappear or get worse. After a concussion, returning to sports and school is a gradual process that should be carefully managed and monitored by a health care professional.
3. Remember: Concussions affect people differently. While most athletes with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

WHY SHOULD AN ATHLETE REPORT THEIR SYMPTOMS?

If an athlete has a concussion, his/her brain needs time to heal. While an athlete's brain is still healing, s/he is much more likely to have another concussion. Repeat concussions can increase the time it takes to recover. In rare cases, repeat concussions in young athletes can result in brain swelling or permanent damage to their brain. They can even be fatal.


Penn High School has partnered with Saint Joseph Sports Medicine Institute to provide concussion baseline testing using the Cogstate Computerized Cognitive Assessment Tool (CCAT) for students in the following contact sports:

Football
Cheerleading
Soccer
Wrestling
Basketball
Baseball (catchers)
Softball (catchers)
Diving
Pole Vault

More information about baseline testing can be found at www.Axonsports.com

The families of student athletes will receive information about testing and be provided with consent forms to fill out from the PHS Athletic Trainers

SAINT JOSEPH
HEALTH SYSTEM

JOIN THE CONVERSATION  www.facebook.com/CDCHeadsUp



TO LEARN MORE GO TO >> WWW.CDC.GOV/CONCUSSION

Content Source: CDC's Heads Up Program. Created through a grant to the CDC Foundation from the National Operating Committee on Standards for Athletic Equipment (NOCSAE).

SUDDEN CARDIAC ARREST

A Fact Sheet for Parents

FACTS

Sudden cardiac arrest is a rare, but tragic event that claims the lives of approximately 500 athletes each year in the United States. Sudden cardiac arrest can affect all levels of athletes, in all sports, and in all age levels. The majority of cardiac arrests are due to congenital (inherited) heart defects. However, sudden cardiac arrest can also occur after a person experiences an illness which has caused an inflammation to the heart or after a direct blow to the chest.

WARNING SIGNS

There may not be any noticeable symptoms before a person experiences loss of consciousness and a full cardiac arrest (no pulse and no breathing).

Warning signs can include a complaint of:

- Chest Discomfort
- Unusual Shortness of Breath
- Racing or Irregular Heartbeat
- Fainting or Passing Out

EMERGENCY SIGNS – Call EMS (911)

If a person experiences any of the following signs, call EMS (911) immediately:

- *If an athlete collapses suddenly during competition*
- *If a blow to the chest from a ball, puck or another player precedes an athlete's complaints of any of the warning signs of sudden cardiac arrest*
- *If an athlete does not look or feel right and you are just not sure*

Developed and Reviewed by the Indiana Department of Education's Sudden Cardiac Arrest Advisory Board
(1-7-15)

How can I help my child prevent a sudden cardiac arrest?

Daily physical activity, proper nutrition, and adequate sleep are all important aspects of life-long health. Additionally, parents can assist student athletes prevent a sudden cardiac arrest by:

- Ensuring your child knows about any family history of sudden cardiac arrest (onset of heart disease in a family member before the age of 50 or a sudden, unexplained death at an early age)
- Ensuring your child has a thorough pre-season screening exam prior to participation in an organized athletic activity
- Asking if your school and the site of competition has an automatic defibrillator (AED) that is close by and properly maintained
- Learning CPR yourself
- Ensuring your child is not using any non-prescribed stimulants or performance enhancing drugs
- Being aware that the inappropriate use of prescription medications or energy drinks can increase risk
- Encouraging your child to be honest and report symptoms of chest discomfort, unusual shortness of breath, racing or irregular heartbeat, or feeling faint

What should I do if I think my child has warning signs that may lead to sudden cardiac arrest?

1. *Tell your child's coach about any previous events or family history*
2. *Keep your child out of play*
3. *Seek medical attention right away*

SUDDEN CARDIAC ARREST

A Fact Sheet for Student Athletes

FACTS

Sudden cardiac arrest can occur even in athletes who are in peak shape. Approximately 500 deaths are attributed to sudden cardiac arrest in athletes each year in the United States. Sudden cardiac arrest can affect all levels of athletes, in all sports, and in all age levels. The majority of cardiac arrests are due to congenital (inherited) heart defects. However, sudden cardiac arrest can also occur after a person experiences an illness which has caused an inflammation to the heart or after a direct blow to the chest. Once a cardiac arrest occurs, there is very little time to save the athlete, so identifying those at risk before the arrest occurs is a key factor in prevention.

WARNING SIGNS

There may not be any noticeable symptoms before a person experiences loss of consciousness and a full cardiac arrest (no pulse and no breathing).

Warning signs can include a complaint of:

- Chest Discomfort
- Unusual Shortness of Breath
- Racing or Irregular Heartbeat
- Fainting or Passing Out

EMERGENCY SIGNS – Call EMS (911)

If a person experiences any of the following signs, call EMS (911) immediately:

- *If an athlete collapses suddenly during competition*
- *If a blow to the chest from a ball, puck or another player precedes an athlete's complaints of any of the warning signs of sudden cardiac arrest*
- *If an athlete does not look or feel right and you are just not sure*

How can I help prevent a sudden cardiac arrest?

Daily physical activity, proper nutrition, and adequate sleep are all important aspects of life-long health. Additionally, you can assist by:

- Knowing if you have a family history of sudden cardiac arrest (onset of heart disease in a family member before the age of 50 or a sudden, unexplained death at an early age)
- Telling your health care provider during your pre-season physical about any unusual symptoms of chest discomfort, shortness of breath, racing or irregular heartbeat, or feeling faint, especially if you feel these symptoms with physical activity
- Taking only prescription drugs that are prescribed to you by your health care provider
- Being aware that the inappropriate use of prescription medications or energy drinks can increase your risk
- Being honest and reporting symptoms of chest discomfort, unusual shortness of breath, racing or irregular heartbeat, or feeling faint

What should I do if I think I am developing warning signs that may lead to sudden cardiac arrest?

1. *Tell an adult – your parent or guardian, your coach, your athletic trainer or your school nurse*
2. *Get checked out by your health care provider*
3. *Take care of your heart*
4. *Remember that the most dangerous thing you can do is to do nothing*